

Measure	CPT/CPT CATEGORY II/HCPCS/ICD-10 Code(s)	Description	Exclusions
Breast Cancer Screening	CPT/HCPCS codes: 77061-77063, 77065 -77067	Indicates whether a female age 52 - 74 had a mammogram done from 27 months prior to the measurement period to the end of the measurement period.	<p>Bilateral mastectomy ICD-10 code: Z90.13 Two unilateral mastectomies ICD-10 codes: Z90.11 AND Z90.12 Unilateral mastectomy with bilateral modifier CPT codes: 19180, 19200, 19220, 19240, 19303-19307</p> <p>Patients who used hospice services anytime during the measurement year.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.</p>
CAD Statin Therapy	At least one high or moderate intensity statin dispensing event during the measurement year.	The percentage of females age 40 - 75 and males age 21 - 75 who were identified as having atherosclerotic cardiovascular disease (ASCVD) and were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year.	<p>Patients with a diagnosis of pregnancy (ICD-10 codes: 0000-09989, 09A*, Z037*, Z331-Z3493, Z36) during the measurement year or year prior. Patients who had in vitro fertilization (HCPCS Procedure Codes: S4015, S4016, S4018, S4020, S4021) performed during the measurement year or year prior. Patients who had a prescription for clomiphene during the measurement year or year prior. Patients with evidence of ESRD (CPT Procedure Codes: 90935, 90937, 90940, 90945, 90947, 90997, 90999, 99512 OR HCPCS Procedure Codes: G0257, S9339 OR ICD-10 codes: M60.80, N18.5, N18.6, Z911.5, Z99.2) during the measurement year or year prior. Patients with a diagnosis of cirrhosis (ICD-10 codes: K70.30, K70.31, K71.7, K74.3-K74.5, K74.60, K74.69, P78.81) during the measurement year or year prior. Patients with diagnosis of myalgia, myositis, myopathy or rhabdomyolysis (ICD-10 codes: G72.0, G72.2, G72.9, M60.80, M62.82, M79.10) during the measurement year or year prior. Patients with hospice services (CPT Procedure Codes: 99377, 99378) during the measurement year.</p> <p>Patients who used hospice services anytime during the measurement year and excludes patients receiving palliative care during the measurement year.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.</p>
Colorectal Cancer Screening	Colorectal cancer screening tests include the following: a fecal occult blood test during the measurement year, a flexible sigmoidoscopy or CT colonography during the measurement year or the previous four years, a stool DNA (sDNA) with FIT test during the measurement year or previous two years, or a colonoscopy during the measurement year or the previous nine years. Fecal Occult Blood Test (FOBT) CPT/HCPCS codes: 82270, 82274, G0328 Flexible Sigmoidoscopy CPT/HCPCS codes: 45330-45335, 45337, 45338, 45340-45342, 45346, 45347, 45349, 45350, G0104 CT Colonography CPT code: 74261-74263 Stool DNA (sDNA) with FIT test CPT/HCPCS code: 81528 Colonoscopy CPT/HCPCS codes: 44388-44394, 44397, 44401-44408, 45355, 45378-45393, 45398, G0105, G0121	Indicates whether a patient, age 45 - 75, had appropriate screening for colorectal cancer.	<p>Colorectal Cancer ICD-10 codes: C18.0-C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048 Total Colectomy CPT codes: 44150-44153, 44155-44158, 44210-44212</p> <p>Patients with hospice service during the measurement year or dispensed dementia meds during the measurement year or year prior.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions. Patients who used hospice services anytime during the measurement year and excludes patients receiving palliative care during the measurement year.</p>

Measure	CPT/CPT CATEGORY II/HCPCS/ICD-10 Code(s)	Description	Exclusions
Controlling High Blood Pressure (BP)	<p>Systolic CPT codes: 3074F, 3075F, 3077F AND Diastolic CPT codes: 3078F, 3079F, 3080F</p> <p>The BP reading must be on or after the visit with the 2nd diagnosis of hypertension.</p>	Patients age 18 - 85 who had a diagnosis of hypertension (HTN) and whose blood pressure was adequately controlled (<140/90) during the measurement year.	<p>Patients with evidence of end-stage renal disease (ESRD) or a kidney transplant on or prior to December 31 of the measurement year. Documentation in the medical record must include a dated note including evidence of ESRD, kidney transplant or dialysis. All patients with a diagnosis of pregnancy during the measurement year.</p> <p>All patients who had a non-acute inpatient stay during the measurement year. Also excludes all patients with hospice services during the measurement year and excludes patients receiving palliative care during the measurement year.</p> <p>Patients who used hospice services anytime during the measurement year.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.</p>
Diabetes Eye Exam	<p>Billed by an optometrist or ophthalmologist CPT codes: 67028, 67030, 67031, 67036, 67039-67043, 67101, 67105, 67107, 67108, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92225-92228, 92230, 92235, 92240, 92250, 92260, 99203-99205, 99213-99215, 99242-99245 HCPCS codes: S0620, S0621, S3000 Unilateral Eye Enucleation CPT codes: 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114 Patient must have two unilateral eye enucleations or one unilateral enucleation with a bilateral modifier.</p> <p>Billed by any provider type during each measurement year. CPT codes: 2022F, 2023F, 2024F, 2025F, 2026F, 2033F, 3072F, 92229</p>	Indicates whether a patient with Type 1 or Type 2 diabetes age 18 - 75 had a dilated retinal eye exam performed by an optometrist or ophthalmologist in the current year OR a dilated retinal eye exam performed by an optometrist or ophthalmologist during the year prior to the measurement year, along with a diagnosis of uncomplicated diabetes. If any other diagnosis showing complicated diabetes is received, the test must be performed annually.	<p>Patients if they had gestational or steroid-induced diabetes during the measurement year AND that patient did not have any other diabetes diagnosis in any setting in the measurement year or year prior. Also excludes anyone with hospice services or palliative care services during the measurement year.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.</p>
Diabetes HbA1c Less Than 9%	<p>CPT codes: 3044F, 3051F and 3052F or LOINC codes: 17856-6, 4548-4, 4549-2 with a Lab Result Numeric Value between 0 and 9</p>	Diabetic patients age 18 - 75 whose HbA1c level is less than 9%.	<p>Patients if they had gestational or steroid-induced diabetes during the measurement year AND that patient did not have any other diabetes diagnosis in any setting in the measurement year or year prior. A diagnosis of HIV (anytime prior to or during the measurement year). Also excludes anyone with hospice services and patients receiving palliative care during the measurement year.</p> <p>Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.</p>
Medication Adherence for Cholesterol	Pharmacy claims	Patients age 18 or older who adhere to their prescribed drug therapy for statin cholesterol medications. Patients fill their prescriptions in the targeted drug class at least 80% or higher of the days prescribed in the measurement year.	<p>Any patient diagnosed with end-stage renal disease (ESRD).</p> <p>Patients who used hospice services anytime during the measurement year.</p>
Medication Adherence for Hypertension	Pharmacy claims	Patients age 18 or older who adhere to their prescribed drug therapy for renin angiotensin system (RAS) antagonists (angiotensin converting enzyme inhibitor (ACEI), angiotensin receptor blocker (ARB) or direct renin inhibitor medications). Qualifying "Blood Pressure medication" includes ACEI, an ARB, or a direct renin inhibitor drug. Patients must fill their prescriptions in the targeted drug class at least 80% or higher of the days prescribed in the measurement year.	<p>Any patient diagnosed with end-stage renal disease (ESRD).</p> <p>Patients who used hospice services anytime during the measurement year.</p>

Measure	CPT/CPT CATEGORY II/HCPCS/ICD-10 Code(s)	Description	Exclusions
Medication Adherence for Diabetes Medications	Pharmacy claims	Patients with Type 1 or Type 2 diabetes age 18 or older who adhere to their prescribed drug therapy across four classes of diabetes medications: biguanides, sulfonylureas, thiazolidinediones, and Dipeptidyl Peptidase (DPP)-IV Inhibitors. Patients must fill their prescriptions in the targeted drug class at least 80% or higher of the days prescribed in the measurement year.	Any patient diagnosed with end-stage renal disease (ESRD). Patients who used hospice services anytime during the measurement year.
Medication Reconciliation Post Discharge	CPT/HCPCS codes: 99483, 99495, 99496, 1111F Measure looks at any inpatient admission to hospital, nursing home, rehab, etc.	Patients 18 and older with an inpatient admission during the measurement year for whom medications were reconciled from the date of discharge through 30 days after discharge (31 days total) by a prescribing practitioner, clinical pharmacist or registered nurse. Only physicians and qualified healthcare providers (MD, DO, NP, etc.) can file TCM codes 99495/99496.	Patients who used hospice services anytime during the measurement year.
Osteoporosis Screening in Older Women	CPT/HCPCS codes: 76977, 77078, 77080, 77081, 77085, 77086	Indicates the the percentage of women 65-85 years of age who received one or more osteoporosis screening tests during the measurement year or the prior year that is on or after the member's 65th birthday.	Women who had a hospice service during the measurement year, frailty or advanced illness claim during the measurement year or one year prior. Women who had an encounter for osteoporosis therapy within 12 months prior to the index event date. Women who had a prescription for an osteoporosis drug within 12 months prior to the index event date. Women who used palliative services anytime during the intake period to the end of the measurement year. Note: See our Advanced Illness and Frailty Guide for coding assistance if a patient may meet criteria for measure exclusions.
Statin Use in Persons with Diabetes	Pharmacy claims	Patients age 40 - 75 who have had two diabetes medication fills and have also received a statin or a statin combination fill in the measurement period.	Any patient diagnosed with End-Stage Renal Disease (ESRD) and any patients enrolled in hospice care at any time during the measurement period. Also, members with the following diagnosis code based conditions coded during the measurement period will be excluded: rhabdomyolysis, myopathy, pregnancy, lactation, liver disease, pre-diabetes and polycystic ovary syndrome. Patients who used hospice services anytime during the measurement year.

These codes are to be used for reference only. This is general information and not a guarantee of payment. Blue Advantage® PPO is provided by Patrius Health, an independent licensee of the Blue Cross and Blue Shield Association. Current Procedural Terminology © 2022 American Medical Association. All rights reserved. Applicable FARS/DFARS apply.