2025 COMMON HCC ICD-10 CODES



When documenting chronic conditions, remember: 1) Coding is based on clear documentation that includes a diagnostic statement and ongoing treatment plan. Any condition that is taken into account or affects patient care, treatment or management at the time of the encounter should be documented and coded. 2) The encounter note should be a face-to-face visit and be complete, legible, concise, and contain the provider signature with credentials, 3) Avoid the use of uncertain diagnoses such as "suggestive of " "suspected" "consistent with" or "probable" and code to the highest level of certainty for the encounter/visit.

| Diabetes Mellitus (DM) | TYPE 1 | TYPE 2 | Alzheimer's disease unspecified | G30.9 |
|--|----------------|-------------------|--|--------------------------------------|
| Diabetes mellitus without complications | E10.9 | E11.9 | Alzheimer's disease: Must be specifically confirmed by the physician to o | |
| Combination Codes: Utilize combination codes that link disease. If there are multiple complications of diabetes m | | | that the Alzheimer's code be paired with the additional F02.8- codes as a Alzheimer's. The physician does not have to mention dementia to code it. | |
| each of the diabetes mellitus combination codes. | | | Epilepsy, unspecified, not intractable, without status epilepticus | G40.909 |
| Diabetes mellitus with hyperglycemia | E10.65 | E11.65 | Unspecified convulsions | R56.9 |
| Hyperglycemia: When hyperglycemia, poorly controlled out of control is documented, diabetes mellitus with hyp The word uncontrolled can mean either hypo- or hyperg | erglycemia sh | nould be coded. | Do not assign R56.9 when a patient has had a seizure disorder or recurrer seizure disorder or recurrent seizures are present utilize appropriate code to | nt seizures. Wher from Category G |
| insufficient documentation to code E10.65/E11.65. | iyoonna, mon | 51010, 11 10 | Cardiovascular Conditions | CODE |
| Diabetes mellitus with diabetic neuropathy, unspecified | E10.40 | E11.40 | Hypertensive heart disease with heart failure | l11.0 |
| Diabetes mellitus with polyneuropathy | E10.42 | E11.42 | Use additional code to identify type of heart failure. | I50 |
| Dishada Badika willa Kida w Osmalisation | TVDE 4 | TVDE 0 | Hypertensive heart disease without heart failure can be coded when there | |
| Diabetes Mellitus with Kidney Complications | TYPE 1 | TYPE 2 | diagnosis and a heart disease code from the range I51.4-I51.7; I51.89; I5 | 1.9. No additiona |
| Diabetes mellitus with diabetic chronic kidney disease | E10.22 | E11.22 | code is needed other than I11.9. | 1100 |
| Diabetes mellitus with diabetic nephropathy | E10.21 | E11.21 | Hypertensive chronic kidney disease with stage 5 chronic kidney disease or end stage renal disease | 112.0 |
| Diabetic chronic kidney disease (CKD) and nephro nephropathy and CKD are documented, code diabetic | | | Use additional code to identify stage of CKD. | N18.5-N18.6 |
| Use additional code to also to identify stage of CKD. | N18.1- N18.6 | N18.1- N18.6 | Hypertensive chronic kidney disease with stage 1 through stage 4 | 112.9 |
| | | | chronic kidney disease, or unspecified chronic kidney disease | |
| Diabetes Mellitus with Ophthalmic Complications | TYPE 1 | TYPE 2 | Use additional code to identify stage of CKD. | N18.1-N18.4 |
| Diabetes mellitus with unspecified diabetic retinopathy | E10.31 | E11.31 | | N18.9 |
| Ophthalmic complications: If adequately documented, code specifically as to type and designate right, left, bilateral or unspecified eye. | E10.3- | E11.3- | Hypertensive heart and chronic kidney disease with heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease | 113.0 |
| | | | Use additional code to identify heart failure type and | I50, N18.1- |
| Diabetes Mellitus with Circulatory Complications | TYPE 1 | TYPE 2 | stage of CKD. | N18.4, N18.9 |
| Diabetic peripheral angiopathy without gangrene | E10.51 | E11.51 | Use additional code to identify stage of CKD. | N18.1-N18.4 N18.9 |
| Diabetic peripheral angiopathy with gangrene | E10.52 | E11.52 | Hypertensive heart and chronic kidney disease without heart failure | 113.11 |
| Diabetes mellitus with foot ulcer | E10.621 | E11.621 | and with stage 5 chronic kidney disease or end stage renal disease | 113.11 |
| Use additional code to identify site of ulcer. | L97.4-, L97. | 5- L97.4-, L97.5- | Use additional code to identify stage of CKD. | N18.5, N18.6 |
| Endocrine, Nutritional and Metabolic Disorders | | CODE | Hypertensive heart and chronic kidney disease with heart failure and | 113.2 |
| Morbid (severe) obesity due to excess calories | | E66.01 | with stage 5 chronic kidney disease or end stage kidney disease | |
| Evaluate for morbid obesity in patients with BMI > or = morbid obesity for patients with comorbidities affected | | evaluate for | Use additional code to identify the type of heart failure and stage of CKD. | I50, N18.5, N18.6 |
| | , , | | Pulmonary hypertension, unspecified | 127.20 |
| Mental, Behavioral and Neurodevelopmental | | CODE | Pulmonary hypertension due to left heart disease | 127.22 |
| Dementia, unspecified without behavioral disturbance | | F03.90- | Pulmonary hypertension due to lung disease and hypoxia | 127.23 |
| Dementia, unspecified with behavioral disturbance | | F03.91- | Cor pulmonale (Chronic) | 127.81 |
| Major Depressive Disorder, single episode, moderate | | F32.1 | Paroxysmal atrial fibrillation | 148.0 |
| Major Depressive Disorder, single episode, severe, with | nout | F32.2 | Longstanding persistent atrial fibrillation | 148.11 |
| psychotic features | 1 11 | F00.0 | Defined as persistent and continuous lasting longer than a year. | |
| Major Depressive Disorder, single episode, severe, with features | 1 psycnotic | F32.3 | Other persistent atrial fibrillation | 148.19 |
| Major Depressive Disorder, single episode, in full remis | esion | F32.5 | Code I48.19 when documented as chronic persistent AF. | |
| Major Depressive Disorder, recurrent, moderate | DOI UTT | F33.1 | Chronic atrial fibrillation, unspecified | 148.20 |
| Major Depressive Disorder, recurrent, severe, without psycl | notic features | F33.2 | Only code chronic AF when documented by the provider. | |
| Major Depressive Disorder, recurrent, severe, without psych | | F33.3 | Permanent atrial fibrillation | 148.21 |
| Document the following components of depression: | one realures | 1 00.0 | Not to be confused with persistent; assign only when permanent AF is o | locumented. |
| 1. Degree (mild, moderate, severe); 2. Episode (single | or recurrent) | ; 3. Status | Unspecified atrial fibrillation | 148.91 |
| (partial or full remission); 4. Presence or absence of partial or full remission) | | | Unspecified atrial flutter | 148.92 |
| Anorexia nervosa, unspecified | | F50.00 | Sick sinus syndrome | 149.5 |

N18.4, N18.9 N18.1-N18.4, N18.9 heart failure 113.11 enal disease N18.5, N18.6 rt failure and 113.2 disease I50.-, N18.5, e and N18.6 127.20 127.22 oxia 127.23 127.81 148.0 148.11 nan a year. 148.19 AF. 148.20 ler. 148.21 manent AF is documented. 148.91 148.92 Anorexia nervosa, unspecified F50.00 Sick sinus syndrome 149.5 Anorexia nervosa, restricting type F50.01 Left ventricular failure, unspecified 150.1 F50.02 Anorexia nervosa, binge eating/purging type Unspecified systolic (congestive) heart failure 150.20 Bulimia nervosa F50.2 Acute systolic (congestive) heart failure 150.21 As of 10/22/2024

PRV20437PAT-2501

| Chronic systolic (congestive) heart failure | 150.22 |
|---|---------|
| Acute on chronic systolic (congestive) heart failure | 150.23 |
| Unspecified diastolic (congestive) heart failure | 150.30 |
| Acute diastolic (congestive) heart failure | 150.31 |
| Chronic diastolic (congestive) heart failure | 150.32 |
| Acute on chronic diastolic (congestive) heart failure | 150.33 |
| Unspecified combined systolic (congestive) and diastolic (congestive) heart failure | 150.40 |
| Acute combined systolic (congestive) and diastolic (congestive) heart failure | 150.41 |
| Chronic combined systolic (congestive) and diastolic (congestive) heart failure | 150.42 |
| Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure | 150.43 |
| Acute right heart failure | 150.811 |
| Acute on chronic right heart failure | 150.813 |
| End stage heart failure | 150.84 |
| Use additional code to identify the type of heart failure as systolic, diastolic or combined, if known (I50.2-I50.43). | |
| Heart failure, unspecified | 150.9 |
| Atherosclerosis of native arteries of extremities with rest pain, right leg | 170.221 |
| Atherosclerosis of native arteries of extremities with rest pain, left leg | 170.222 |
| Atherosclerosis of native arteries of extremities with rest pain, bilateral legs | 170.223 |
| Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, right leg | 170.321 |
| Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, left leg | 170.322 |
| | |

| Diseases of the Respiratory System | CODE |
|--|---------------|
| Unspecified chronic bronchitis | J42 |
| Emphysema, unspecified | J43.9 |
| Chronic obstructive pulmonary disease with (acute) lower respiratory infection | J44.0 |
| Use additional code to identify the infection. | |
| Chronic obstructive pulmonary disease with (acute) exacerbation | J44.1 |
| If COPD with acute exacerbation and COPD with lower respiratory present: document and code both. | infection are |
| Chronic obstructive pulmonary disease, unspecified | J44.9 |
| Severe persistent asthma, uncomplicated | J45.50 |
| Severe persistent asthma with (acute) exacerbation | J45.51 |
| Severe persistent asthma with status asthmaticus | J45.52 |
| Respiratory failure | J96.1-J96.92 |
| Diseases of the Directive System | CODE |

| Diseases of the Digestive System | CODE |
|----------------------------------|--------|
| Crohn's Disease, unspecified | K50.9- |
| Ulcerative colitis, unspecified | K51.9- |

| Diseases of the Musculoskeletal System and Connective Tissue | CODE |
|---|-------------|
| Rheumatoid arthritis, unspecified | M06.9 |
| Rheumatoid arthritis: If known, code specifically regarding with or without rheumatoid factor. Also code specific site as documented. | M05-M06.8A |
| Systemic lupus erythematosus, unspecified | M32.9 |
| Systemic lupus erythematosus (SLE): Documentation must confirm and specify Lupus as systemic to assign a code from category M32. | M32.0-M32.9 |

Genitourinary Conditions

Two separate eGFR values are required to change the CKD stage. When both acute kidney failure and CKD are present, code both conditions.

Chronic kidney disease, stage 3 unspecified

N18.30

| kidney failure and GKD are present, code both conditions. | | | | | |
|---|---|--------|--|--|--|
| | Chronic kidney disease, stage 3 unspecified | N18.30 | | | |
| | Chronic kidney disease, stage 3a | N18.31 | | | |
| | Chronic kidney disease, stage 3b | N18.32 | | | |
| | Chronic kidney disease, stage 4 (severe) | N18.4 | | | |
| | Chronic kidney disease, stage 5 | N18.5 | | | |
| | End stage renal disease | N18.6 | | | |

| Factors Influencing Health Status | CODE | | |
|-----------------------------------|--------------|--|--|
| Body mass index (BMI) | Z68.1-Z68.45 | | |

BMI: Reported BMI >40 requires a weight diagnosis to be documented and coded, such as "Morbid Obesity" to rise to an HCC.

| Long- | term | (cu | rrer | nt) ı | use | of | ins | sulin | l | | | Z7 | '9.4 | |
|-------|------|-----|------|-------|-----|----|-----|-------|---|--|--|------|------|--|
| 16.11 | | | - | - | - | | | | | | | | | |

If the patient is treated with both oral hypoglycemic drugs and insulin, code both Z79.84 and Z79.4.

| Personal history of malignant neoplasm (code specific history of | Z85 |
|--|-----|
| cancer code) | |

Differentiating between "active" and "history of" cancer: When determining active versus history of cancer, look for active treatment, such as chemotherapy, radiation, hormone therapy, watchful waiting, or patient refusal of treatment. Remember, when coding active cancer, document specific primary sites and metastasis (secondary), if present.

| Personal history of transient ischemic attack (TIA)/cerebrovascular | Z86.73 |
|---|--------|
| accident (CVA) without residual effects | |

TIA/CVA: Use history of codes unless patient is having active symptoms of TIA or CVA during the visit. If the patient has residual of a CVA, code the residual symptoms or sequelae to the highest level of specificity I69.3- (i.e., monoplegia, hemiplegia, hemiplegia, hemiparesis, etc.).

| Acquired absence of toe(s), foot and ankle | Z89.4-Z89.449 |
|--|---------------|
| Acquired absence of leg below knee | Z89.5-Z89.519 |
| Acquired absence of leg above knee | Z89.6-Z89.619 |
| Tracheostomy status | Z93.0 |
| Gastrostomy status | Z93.1 |
| lleostomy status | Z93.2 |
| Colostomy status | Z93.3 |
| Cystostomy status | Z93.5- Z93.59 |
| Artificial opening status, unspecified | Z93.9 |
| Heart transplant status | Z94.1 |
| Lung transplant status | Z94.2 |
| Heart and lung transplant status | Z94.3 |
| Liver transplant status | Z94.4 |
| Bone marrow transplant status | Z94.81 |
| Intestine transplant status | Z94.82 |
| Pancreas transplant status | Z94.83 |
| Stem cells transplant status | Z94.84 |
| Presence of fully implantable artificial heart | Z95.812 |

When your patient is immunocompromised while taking anti-rejection medication, consider documenting and coding D84.821 "Immunodeficiency due to drugs."

Dependence on renal dialysis

Z99.2

ICD-10 is the 10th revision of the International Statistical Classification of Diseases and Related Health Problems (ICD), a medical classification list by the World Health Organization (WHO).



An Independent Licensee of the Blue Cross and Blue Shield Association